

## PRECAUTIONARY STATEMENTS

### HAZARD TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Keep away from humans, domestic animals, and pets. If swallowed by humans, domestic animals, or pets this material may reduce the clotting ability of the blood and cause bleeding. Wash thoroughly with soap and water after handling and before eating or smoking.

Note to Physician: If ingested, administer Vitamin K1 intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

**ENVIRONMENTAL HAZARDS:** This product is toxic to fish and wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below mean high water mark.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in original container in a cool, dry place inaccessible to children and pets.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill.

## DO NOT CONTAMINATE FEED OR FOODSTUFFS

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

## RESTRICTED USE PESTICIDE

For retail sale to and only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certificate.

## FOR DISTRIBUTION AND USE ONLY IN MONTANA

This label must be in the possession of the user at the time of applications. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

# rozol®

## Ground Squirrel Oat Bait

Active Ingredient: chlorophacinone . . . . 0.005%  
Inert Ingredients . . . . . 99.995%  
Total . . . . . 100.000%

## KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements.

**LIPHA**TECH®

3600 W. Elm Street  
Milwaukee, WI 53209  
800-351-1476

NET CONTENTS: 40 LBS.

## DIRECTIONS FOR USE

For use only against Columbian and Richardson ground squirrels (*Spermophilus columbianus* and *S. richardsonii*). Do not apply on roads, near residential or public areas or over water.

**Hand or Spot Baiting:** Apply only at a time when ground squirrels are readily accepting grains. Attempt to time bait application during a period when rain is not expected for four (4) days after the first application. A light shower will unlikely affect bait, but a soaking rain will reduce the efficacy of the bait. This product must be eaten by the target squirrels more than once within 3 to 4 days to be effective. A single application or a single application at double the application rate will result in poor control. Apply only at the frequency and rate recommended to obtain the best results and to present the least hazard to nontarget animals. Do not exceed application rate. Do not place bait on soft mound dirt or down the burrow opening. Do not pile bait. Do not broadcast bait by mechanical spreaders or other methods. Apply bait only to bare ground around ground squirrel burrows in non-crop areas, rangeland, pasture, alfalfa, wheat, oats and barley. Remove livestock from the treatment area until the bait is consumed by the squirrels to prevent bait consumption or trampling by livestock.

**Pre-treatment:** Do not apply bait unless tests with untreated oats show that untreated oats are completely consumed by the target squirrels within two (2) days.

**Treatment:** Use a calibrated measurer or dipper (such as a 1 inch copper pipe cap attached to a 12 inch length of 1/4 inch rod) to apply one tablespoon (0.3 oz. or 9 grams) of bait per active burrow. Scatter bait evenly on bare ground near each burrow entrance. Make a second application in the same manner four days after the first application.

**Bait station placements:** Apply bait in tamper-resistant bait stations where it will be readily accessible to ground squirrels, preferably near their burrows. Assure that stations cannot be opened or disturbed by children, domestic animals or non-target wildlife. Stations must be constructed to permit entry by target species but not by larger animals. Construct bait stations to inhibit spillage of bait. Secure bait stations so that wind, livestock, pets and wildlife cannot overturn them. Check bait stations frequently to insure that an uninterrupted supply of bait is available. Maintain an uninterrupted supply of bait for at least 30 days or until all signs of feeding have stopped. Bait stations may be maintained on a permanent basis to prevent reinfestation. Bait stations should be located convenient to squirrel burrows and should be placed 20 to 100 feet apart, depending on population density.

**Post-treatment:** Search treatment area for squirrel carcasses. Place any carcasses found, individually, down ground squirrel burrows to reduce hazards to nontarget animals that may consume the carcasses.